

Factors responsible for productivity of food legumes: Findings of a farmer survey in the Kurunegala district, Sri Lanka

Short communication

M.P. Hettiarachchi^{1*}, W.A.J.M. De Costa^{1**} and S.J.B.A. Jayasekera²

¹Department of Crop Science, Faculty of Agriculture, University of Peradeniya, Sri Lanka.

²Department of Horticulture, Wayamba Campus, Rajarata University of Sri Lanka, Makandura, Sri Lanka.

Accepted 10 November 1998

ABSTRACT

Food legumes are an important component in upland cropping systems of subsistence farmers in Sri Lanka. The objective of this study was to understand the factors responsible for the gap between potential yield and farmer yield in Kurunegala district. Results revealed that information on improved varieties of cowpea (*Vigna unguiculata* (L.) Walp), greengram (*Vigna radiata* (L.) Wilczek) and groundnut (*Arachis hypogaea* L.) and on better management practices have not reached the farmers.

Key words: Food legumes, productivity, Sri Lanka, yield.

Food legumes are an alternative source of protein for the majority of the population in Sri Lanka (Gunaseena and Herath 1987). In 1995, cowpea (*Vigna unguiculata* (L.) Walp), greengram (*Vigna radiata* (L.) Wilczek), soybean (*Glycine max* (L.) Merrill), groundnut (*Arachis hypogaea* L.) and common bean (*Phaseolus vulgaris* L.) were cultivated in an area of 60,902 ha to produce 79,475 mt of yield (Anon. 1995a). In Sri Lanka, Kurunegala (longitude - 80° 30'E, latitude - 70° 30'N, mean annual rainfall 1800mm, mean temperature 30°C, soil-Reddish Brown Earth (Rhodustalfs), Low Humic Gley (Tropaqualfs) and Non Calcic Brown (Haplustalfs) is one of the major legume growing districts, especially of greengram (24.5%), cowpea (21.3%) and groundnut (12%) (Anon. 1995b).

The average yields of greengram (0.35 t ha⁻¹), cowpea (0.32 t ha⁻¹) and groundnut (0.55 t ha⁻¹) obtained by farmers in Kurunegala district are much lower than the potential yields (greengram- 0.88 t ha⁻¹, cowpea- 0.82 t ha⁻¹ & groundnut- 0.97 t ha⁻¹) obtainable from farmer fields (Jayasekera and Hettiarachchi 1987; Jayasekera and Ariyaratne 1988). The objective of the present paper is to provide specific information on cowpea, greengram

and groundnut production in Kurunegala district and to highlight problems and constraints of legume cultivation from the farmer's viewpoint.

One hundred farmers were selected using the stratified random sampling technique. Farmers were selected in the survey to represent the larger legume growing villages: Yakadapotha, Meegallewa, Paluwaduressa, Mahagirilla, Udagirilla, Waduressa and Madapotha in the Kotawehera range and Palliyakale, Boraluwewa, Udawewa, Korakahawetiya and Hirigolla in the Kobeigane range. A structured questionnaire was used to collect cultivation information on greengram, cowpea and groundnut from farmers in the selected areas.

Extent and purpose of legume cultivation in the survey areas

In the Kotawehera area, all the surveyed farm families cultivated greengram and cowpea during southwest monsoon (May - July) and northeast monsoon (October -January) seasons in the uplands. Groundnut was cultivated only during southwest monsoon by 80% of the farmers surveyed. In Kobeigane, only 50% of the farmers surveyed cultivated legumes. In this area, cultivation was confined to lowlands during southwest monsoon because there was not enough irrigation water to grow a rice crop. Here, the uplands were not used for legume cultivation because of lack of irrigation facilities.

*Present address: Department of Crop Science, Faculty of Agriculture, University of Ruhuna, Kamburupitiya, Sri Lanka.

** Corresponding author

Cultivated varieties and sources of seeds

Table 1 shows that in all three species of legumes, a majority of farmers surveyed still cultivate the older varieties and the adoption of recently-released varieties is low. Local varieties of greengram are known among the farmers as *Pinna mung* and *Alu mung* whereas *MI 5* is also known as *Tel mung*. Among the cowpea varieties, Arlington is cultivated during both seasons whereas *MI 35*, *Vijaya* and *Varuni* are cultivated only during the southwest monsoon season. The survey also revealed that the majority of farmers used their own seeds (Table 1) because they were confident about the seed viability. As a result of the continued use over several seasons (Table 2), the genetic purity of many varieties have been significantly reduced and often the improved varieties are mixed with local varieties. Even when the farmers purchase their seeds, only a minority uses the sales centres of the Department of Agriculture (DOA).

Table 1. Cultivated varieties, sources of seeds and some cultural practices of three selected legumes in Kurunegala district, Sri Lanka.

Factor	Percentage of farmer surveyed		
	Greengram	Cowpea	Groundnut
Varieties grown			
Old varieties	Local varieties - 50	Arlington - 100	Red spanish & V87 - 100
New varieties	MI 5 - 75 Harsha - 15	MI 35 - 50 Vijaya & Varuni - 25	Tissa & Walawe - 20
Sources of seeds			
Own seeds used	80*	80	90
DOA sales	30	20	10
Private traders	70	80	90
Sowing - 2 seeds/hill - 3 seeds/hill	100 -	50 50	100 -
Fertilizer application			
Northeast monsoon season	85	70	-
Southwest monsoon season	40	10	20
Crop duration	2-3 months - 40 3-3 1/2 months - 60	3 months - 100	3-3 1/2 months - 100
Harvesting	One pick - 10 3-4 picks - 70 7-8 picks - 20	3-4 picks - 60 5-6 picks - 40	One pick - 100

*DOA - Department of Agriculture, Sri Lanka

*Farmers use seeds from all sources in same season.

Cultural practices

Table 1 shows some of the key cultural practices and the percentages of surveyed farmers who use them. Farmers predominantly used two seeds per hill during sowing. Although it was advantageous to use more than two seeds per hill to ensure adequate crop establishment under adverse soil conditions, higher seed cost, prevented the farmers from using a high seed rate. Fertilizer rates applied were generally higher during the northeast monsoon season. The greater probability of success of the northeast

monsoon crop encouraged the farmers to use more inputs. Frequent failures of southwest monsoon crop, mainly due to drought, discouraged farmer investment in costly inputs such as fertilizer.

Among species surveyed, greengram had varieties with a wider range of maturity. This was a favourable characteristic in this environment where drought is a major cause of crop failure. Both greengram and cowpea allowed several picks of harvest. This trait also added flexibility to the crop by allowing an early harvest before an environmental stress sets in or allowing recovery of yield following a stress period. Both the above characters were major reasons for higher farmer preference for greengram and cowpea in the Kurunegala district.

Pest and disease problems

In both the areas surveyed, legume yields were severely reduced due to pest and disease infestation. The most common pests and diseases in the selected legumes are *Maruca testulalis* and *Lampedes boeticus* (80%), *Ophiomyia phaseoli* (60%), *Callosobruchus chinensis* (60%), *Aphis sp.* & Thrips (40%), Rodents (40%), Yellow Mosaic Virus (70%), fungal diseases (50%) and Wilt (25%). Only 30% of the surveyed farmers applied commercial fungicides to their crops whereas 20% applied sulphur as a treatment. On the other hand, of the farmers surveyed 40% used pesticides. This also highlighted the reluctance of the farmers to invest in a greater level of inputs. A majority of farmers felt that pesticide application was not economical because of marginal increase in yield. Another reason for the low use of pesticides was their non-availability at the required times.

More than 40% of farmers who used pesticides did not know the names of pesticides used and very often a chemical for a given pest or disease was recommended to them by the trader from whom it was bought.

Yields and selling prices

According to the farmers surveyed, only greengram and cowpea gave appreciable yields whereas the groundnut yields were minimal. Greengram and cowpea yields of all farmers surveyed (Table 1) were lower than the potential yield of the varieties (1600 & 1800 kg/ha respectively-Department of Census & Statistics). Only 30% and 20% of the surveyed farmers obtained yields greater than 500 kg/ha for greengram and cowpea respectively.

The selling price was dependent on the specific varietal characteristics. For example, the smaller

Table 2. Farmers' perception of factors responsible for low yields, period of cultivating the same variety and awareness of new varieties of selected legumes in Kurunegala district, Sri Lanka.

Factors	Percentage of farmers surveyed
Reasons for low yield	
Drought	100
Water-logging due to high rainfall (in northeast monsoon)	30
Not using newly-released high-yielding varieties	80
Crop damage due to rabbits, wild boars, rats and cattle	60
Low soil fertility	50
Lack of fertilizer application	50
Increased pest and disease problems due to delayed and staggered cultivation	30
Low levels of input use and crop management due to financial constraints	80
Periods of cultivations of same variety	
2-3 years	40
5-6 years	20
13-14 years	20
20-25 years	20
Awareness of new varieties	
Greengram	30
Cowpea	20
Groundnut	10

seeds of the new greengram variety *Harsha* fetched a lower price in Kurunegala district whereas *MI 5* always fetched a higher price. Hence, even the farmers who knew about *Harsha* did not want to grow it because of the ensuing selling problems.

Factors responsible for low yields

Table 2 lists the factors responsible for low legume yields as perceived by the surveyed farmers. Drought and low levels of input use and crop management were identified by a majority of surveyed farmers as factors causing low yields. Among the inputs, failure to use newly-released high-yielding varieties was perceived as a major factor causing low yields. Out of the surveyed farmers, 70%, 80% and 90% mentioned that they did not receive information on newly-released varieties of greengram, cowpea and groundnut respectively. Specifically, only 15% knew about *Harsha*, the newly-introduced greengram variety and this was only through the Upland Development Project conducted through the North-Central Provincial Council. The new cowpea varieties *Vijaya* and *Varuni* were known to only 25% of the surveyed farmers which was also through the above project.

Outlook for the future

In spite of the low yields and constraints described above, all (*i.e.* 100%) the interviewed farmers preferred to continue cultivation of legumes because of several reasons: (a) Legumes need a lower level of inputs and crop management as compared to other high-value upland cash crops such as chilli or

onions; (b) Water requirement of legumes is lower than for alternative upland crops; (c) Legumes can be used either for selling or domestic consumption as a major food item in the daily diet, unlike crops such as chilli or onions. On the whole, the farmers felt that under the prevailing environmental and socio-economic conditions in the region, there was no choice but to cultivate legumes as their major upland annual crop.

The present study identified the salient features of legume cultivation and highlighted the factors responsible for lower farmer yields of greengram, cowpea and groundnut in the Kurunegala district. Although the survey was done within a limited area of the Kurunegala district, the agro-climatic and socio-economic environment that prevailed in the survey area was representative of many of the legume-growing areas in the rainfed dry and intermediate zones of Sri Lanka (Herath and Suraweera 1987).

One finding of the survey was the lack of flow of information about newly-released improved legume varieties to the farmers who continue to grow local varieties of low purity which are mostly produced by them. This would mean that all the research effort in producing new legume varieties would be wasted. This gap in the flow of information about new technology has to be addressed as quickly as possible by strengthening the extension service and its linkages with the research centres and the farmers.

It was also revealed that in addition to the lack of knowledge about new varieties, there were no channels through which farmers could get precise information about correct cultural practices and pest and disease control. Both extension services and infra-structure services such as sales centres for seeds and other inputs need to be improved to uplift the general level of productivity and farmer income in legume cultivation. This would help to break the dependence of most of these farmers on private traders to whom the farmers are bound to sell their produce at lower prices. In addition to the extension services, farmer training and education should be strengthened to make the farmers aware of the changing market trends and how best to adjust to them in order to maximize their profit.

ACKNOWLEDGEMENTS

Authors wish to thank Agricultural Officers in Provincial Agricultural Director's Office at Kurunegala, Agricultural Instructors in Kotawehera and Kobeigane Agrarian Services Centres and farmers in these areas for providing information on legume cultivation.

REFERENCES

- Anonymous 1995a Annual report, Dept. of Census & Statistics, Colombo, Sri Lanka.
- Anonymous 1995b Socio-Economic and Planning Centre, Department of Agriculture, Peradeniya, Sri Lanka.
- Gunasena HPM and Herath HMG (eds.) 1987 Improved Production and Utilization of Food Legumes in Sri Lanka, Proceedings of Symposium, Peradeniya, Sri Lanka.
- Herath HMG and Suraweera DEF 1987 Socio-economic aspects of food legume production in Sri Lanka. In: H.P.M. Gunasena and H.M.G. Herath. (eds.) Improved Production and Utilization of Food Legumes in Sri Lanka. Peradeniya, Sri Lanka. pp. 7-22.
- Jayasekera SJBA and Ariyaratne HP 1988 Current status of mung bean improvement for the farming systems in Sri Lanka. In: S. Shanmugasundaram and B.T. McLean. (eds.) Mungbean, Proc. Second International Symp., Asian Vegetable Research and Development Centre, Shanhua, Taiwan. pp. 641-650.
- Jayasekera SJBA and Hettiarachchi K 1987 Cowpea and groundnut improvement programme in Sri Lanka. In: H.P.M. Gunasena and H.M.G. Herath. (eds.) Improved Production and Utilization of Food Legumes in Sri Lanka, Peradeniya, Sri Lanka. pp. 134-146.

EDITORIAL POLICIES

Submission of manuscripts:

The journal publishes research results of broad practical significance for tropical and subtropical agriculture. The subject areas covered include crop genetics, resource agronomy, crop improvement, crop physiology, plant protection, weed science, ecology and sustainable management of tropical and subtropical agricultural environments, restoration of degraded environments, agro-forestry, utilization and domestication of under-exploited plant and animal resources, bio-technological applications, horticulture, perennial crops, animal science including fisheries, food science, post-harvest technology, farming systems, agricultural economics, extension and education.

The papers submitted to the journal should not have been published previously in the same or any other form or language, or being considered for publication elsewhere. However, publications appearing as abstracts or as part of a published lecture or thesis may be submitted as a research paper.

The journal publishes review articles on critical evaluations of existing knowledge in a specialized field, research papers and short communications. The journal favors concise, readable publications relating to research on indigenous under-exploited plant and animal resources, traditional methods and techniques, work containing new methodological approaches, etc.

Language of publication: English

References and editing:

All material submitted for publication will be reviewed. The papers will be edited if necessary to improve clarity and ease of communication. Papers conforming to the conventions of the journal with regard to symbols, illustrations, tables, references, nomenclature, etc. will be processed expeditiously.

Frequency of publication:

The journal is a half-yearly journal with two issues per volume published in the months of June and December.

MANUSCRIPTS

The original and two copies of the manuscript should be submitted.

Length:

Although there is no maximum length prescribed for Review Articles or Research Papers, the authors are requested to write clearly and concisely. Short Communications, which should carry an abstract of about 50 words, should not exceed 1000 words including a maximum of two tables. Long word articles should be structured as prescribed below. Research Papers: Nevertheless, the text should include the objectives of the study, materials and methods used, results obtained, discussion of the results, conclusions and relevant references. Longer articles and review proceedings of relevant conferences and symposia may be published from time to time as supplementary issues of the journal.